arXiv:1005.3761v2 [math.PR] 19 Jan 2012

Random matrix models of stochastic integral type for free infinitely divisible distributions

J. Armando Domínguez Molina* and Alfonso Rocha Arteaga†
Facultad de Ciencias Físico-Matemáticas
Universidad Autónoma de Sinaloa, México

Abstract

The Bercovici-Pata bijection maps the set of classical infinitely divisible distributions to the set of free infinitely divisible distributions. The purpose of this work is to study random matrix models for free infinitely divisible distributions under this bijection. First, we find a specific form of the polar decomposition for the Lévy measures of the random matrix models considered in Benaych-Georges [6] who introduced the models through their measures. Second, random matrix models for free infinitely divisible distributions are built consisting of infinitely divisible matrix stochastic integrals whenever their corresponding classical infinitely divisible distributions admit stochastic integral representations. These random matrix models are realizations of random matrices given by stochastic integrals with respect to matrix-valued Lévy processes. Examples of these random matrix models for several classes of free infinitely divisible distributions are given. In particular, it is shown that any free selfdecomposable infinitely divisible distribution has a random matrix model of Ornstein-Uhlenbeck type $\int_0^\infty e^{-t} d\Psi_t^d$, $d \geq 1$, where Ψ_t^d is a $d \times d$ matrix-valued Lévy process satisfying an $I_{\log}$ condition.

1 Introduction

An ensemble of random matrices is a sequence $(M_d)_{d\geq1}$ where M_d is a $d \times d$ matrix whose entries are random variables. The empirical spectral distribution of M_d is the uniform distribution of its spectrum $\lambda_1, \lambda_2, ..., \lambda_d$, that is the (random) probability measure μ_{M_d} defined as $\mu_{M_d} = \frac{1}{d}\sum_{i=1}^d \delta_{\lambda_i}$. A random matrix model for a probability measure μ is an ensemble $(M_d)_{d\geq1}$ for which the empirical spectral distribution μ_{M_d} converges weakly to μ.

Bercovici and Pata [7] introduced a bijection Λ from the set of classical infinitely divisible distributions to the set of free infinitely divisible distributions to study relations between

---

*jadguez@uas.uasnet.mx
†arteaga@uas.uasnet.mx

classical and free infinitely divisible aspects. Under this bijection Benaych-Georges [6] and Cavanal-Duvillard [8] construct for any classical one-dimensional infinitely divisible distribution μ a random matrix model for the corresponding free infinitely divisible distribution $\Lambda(\mu)$. This include the Wigner and Marchenko-Pastur results, which provide random matrix models of Gaussian Unitary Ensembles and Wishart random matrices for the semicircle and Marchenko-Pastur distributions, respectively.

Specifically, it is shown in [6] that for any infinitely divisible distribution μ on $\mathbb{R}$ there is a random matrix model of Hermitian matrices $(M_d)_{d\geq 1}$ for $\Lambda(\mu)$. Moreover, for each $d \geq 1$ the Fourier transform of M_d is given by

$$\mathbb{E}[\exp(i\operatorname{tr}(M_d A))] = \exp\left[d\mathbb{E}_u \mathcal{C}_\mu\left(\langle u, Au\rangle\right)\right] \;\; A \text{ Hermitian}, \tag{1}$$

where $\mathcal{C}_\mu$ is the cumulant transform of μ, u is a uniformly distributed column random vector in the unit sphere of $\mathbb{C}^d$ and $\langle\cdot,\cdot\rangle$ is the usual Hermitian product of $\mathbb{C}^d$.

Any $\mathbb{R}$-valued Lévy process $\left\{X_t^{(\mu)} : t \geq 0\right\}$ with law μ at time $t = 1$ has associated an $\mathbb{R}$-valued independently scattered random measure. The stochastic integral of a real-valued function h on $[0,\infty)$ with respect to $X_t^{(\mu)}$ written as

$$\int_0^\infty h(t)dX_t^{(\mu)}, \tag{2}$$

is an $\mathbb{R}$-valued infinitely divisible random variable defined in the sense of integrals of non random functions with respect to scattered random measures, see Urbanik and Woyczynski [21] and Rajput and Rosiński [15]; and Sato [19] for the $\mathbb{R}^d$ case. Several important classes of infinitely divisible distributions having this stochastic integral representation in law have been studied recently, see [4], [2], [1], [3] and [9].

Let $I_{\log}(\mathbb{R})$ be the class infinitely divisible distributions μ on $\mathbb{R}$ whose Lévy measures ν_μ satisfy the condition $\int_{|x|>2}\log|x|\,\nu_\mu(dx) < \infty$. It is shown in Jurek and Vervaat [11], Sato and Yamazato [20] and Sato [18], that the class of selfdecomposable distributions on $\mathbb{R}$ is characterized by the stochastic integrals of the form (2) where $h(t) = e^{-t}$ and $\mu \in I_{\log}(\mathbb{R})$ in the following sense. For any $\mu \in I_{\log}(\mathbb{R})$ there exists a selfdecomposable distribution $\widetilde{\mu}$ such that

$$\widetilde{\mu} = \mathcal{L}\left(\int_0^\infty e^{-t}dX_t^{(\mu)}\right) \tag{3}$$

and vice versa, to any selfdecomposable distribution $\widetilde{\mu}$ corresponds a distribution μ in $I_{\log}(\mathbb{R})$ such that (3) holds. This characterization of selfdecomposable distributions as stochastic integrals is related to Ornstein-Uhlenbeck type processes through the Langevin equation. The Langevin equation $dY_t = dX_t^{(\mu)} - Y_t dt$ has stationary solution $\{Y_t : t \geq 0\}$ if and only if $\mu \in I_{\log}(\mathbb{R})$. This stationary solution $\{Y_t\}$ is unique and $\mathcal{L}(Y_t) = \mathcal{L}\left(\int_0^\infty e^{-t}dX_t^{(\mu)}\right)$ for all $t \geq 0$. The process $\{Y_t\}$ is called the stationary Ornstein-Uhlenbeck type process, see Sato [18] and Rocha-Arteaga and Sato [16].

In this work we are concerned with random matrix models for free infinitely divisible distributions corresponding to the image Λ of classical infinitely divisible distributions. We show that for every classical one-dimensional infinitely divisible distribution representable as the stochastic integral (2) there exists a random matrix model for the corresponding free infinitely divisible distribution, consisting of a realization as matrix stochastic integral similar to (2) with respect to an appropriate matrix-valued Lévy process. In particular, the free selfdecomposable distribution $\Lambda(\widetilde{\mu})$ corresponding to the classical selfdecomposable distribution $\widetilde{\mu}$ with stochastic integral representation (3) where $\mu \in I_{\log}(\mathbb{R})$, has a realization as random matrix model of Ornstein-Uhlenbeck type $\left(\int_0^\infty e^{-t} d\Psi_t^{\mu,d}\right)_{d\geq 1}$ where $\Psi_t^{\mu,d}$ is a $d \times d$ matrix-valued Lévy process satisfying an $I_{\log}$-condition. Recently, Pérez-Abreu and Sakuma [14] studied random matrix models for free infinitely divisible distributions as matrix stochastic integrals of Wiener-Gamma type. They considered free infinitely divisible distributions corresponding to the image Λ of classical generalized Gamma convolutions distributions with the so called Wiener-Gamma representation, a type of stochastic integral representation (2) with respect to the Gamma process.

This paper is organized as follows. In Section 2 we find the polar decomposition for the Lévy measures of the Hermitian matrices of the random matrix models (1). In Section 3 we construct random matrix models for free infinitely divisible distributions as realizations of classical matrix stochastic integrals with respect to matrix-valued Lévy processes. We prove that these matrix stochastic integrals have the Fourier transform (1). In Section 4 we provide examples of random matrix models of matrix stochastic integrals for several classes of free infinitely divisible distributions. In particular, the class of free selfdecomposable distributions corresponding to the image Λ of the class of selfdecomposable distributions in (3), has random matrix models of Ornstein-Uhlenbeck type matrix integrals.

2 Polar decomposition of Lévy measures of certain random matrix models

Let $\mathbb{M}_d = \mathbb{M}_{d\times d}(\mathbb{C})$ denote the linear space of $d \times d$ Hermitian matrices with scalar product $\langle A, B\rangle = \operatorname{tr}(AB^*)$ and norm $\|A\| = [\operatorname{tr}(AA^*)]^{1/2}$ where tr denotes trace. Let $\tilde{\mathbb{S}}_{\mathbb{M}_d} = \{V \in \mathbb{M}_d : \operatorname{rank}(V) = 1, \|V\| = 1\}$ the set of Hermitian matrices of rank 1 on the unit sphere of $\mathbb{M}_d$ and let $\tilde{\mathbb{S}}^+_{\mathbb{M}_d} = \{V > 0 : \operatorname{rank}(V) = 1, \|V\| = 1\}$ the set of positive definite matrices of rank 1 on the unit sphere of $\mathbb{M}_d$.

We recall the polar decomposition of Lévy measures on $\mathbb{R}$, see [17] and [4]. The Lévy measure ν of an infinitely divisible distribution μ on $\mathbb{R}$ with $0 < \nu(\mathbb{R}) \leq \infty$, can be expressed as

$$\nu(B) = \int_S \lambda(d\xi) \int_0^\infty 1_B(r\xi)\nu_\xi(dr), \tag{4}$$

where λ is a measure on the unit sphere $S = \{-1, 1\}$ of $\mathbb{R}$ such that $0 < \lambda(S) \leq \infty$ and ν_ξ is a measure on $(0, \infty)$ for each $\xi \in S$ such that $0 < \nu_\xi((0, \infty)) \leq \infty$. Here λ and ν_ξ are called the spherical and radial components of ν, respectively.

Let $\widehat{\mu}$ and $\mathcal{C}_\mu$ denote the Fourier transform and the cumulant transform of an infinitely divisible distribution μ on $\mathbb{R}$, respectively. That is, $\mathcal{C}_\mu$ is the unique continuous function from $\mathbb{R}$ into $\mathbb{C}$ such that $\mathcal{C}_\mu(0) = 0$ and $\widehat{\mu}(z) = \exp(\mathcal{C}_\mu(z))$ for every $z \in \mathbb{R}$.

In [6, Theorem 6.1] it is established that for any infinitely divisible distribution μ on $\mathbb{R}$ there exists a random matrix model of Hermitian matrices $(M_d)_{d\geq 1}$ for $\Lambda(\mu)$. Moreover, in [6, Theorem 3.1] the Fourier transform of M_d, for each $d \geq 1$, is given by

$$\mathbb{E}[\exp(i\mathrm{tr}(M_d A))] = \exp\left[d\mathbb{E}_u \mathcal{C}_\mu(\langle u, Au\rangle)\right] \quad A \text{ Hermitian} \tag{5}$$

where $\mathcal{C}_\mu$ is the cumulant transform of μ, u is a uniformly distributed column random vector in the unit sphere of $\mathbb{C}^d$ and $\langle\cdot,\cdot\rangle$ is the usual Hermitian product of $\mathbb{C}^d$. In the sequel we denote by ω_d the probability measure on the set of matrices of rank 1 induced by the transformation

$$u \to V = uu^*. \tag{6}$$

In the following result we find the specific form of the polar decomposition for the Lévy measures of the random matrix models considered in [6].

Theorem 1 *Let μ be an infinitely divisible distribution on $\mathbb{R}$ with Lévy measure ν and let $(M_d)_{d\geq 1}$ a random matrix model for $\Lambda(\mu)$ where M_d has the Fourier transform (5). Then the Lévy measure ν_{M_d} of M_d is expressed as*

$$\nu_{M_d}(B) = d\int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}} \int_0^\infty 1_B(rV)\,\nu_V(dr)\,\Pi(dV) \quad B \in \mathcal{B}(\mathbb{M}_d \backslash \{0\}),$$

where $\nu_V = \nu^+$ or ν^- according to $V > 0$ or $V < 0$ and $\Pi(dV)$ is a measure on $\tilde{\mathbb{S}}_{\mathbb{M}_d}$ such that

$$\int\limits_{\tilde{\mathbb{S}}_{\mathbb{M}_d}} 1_D(V)\,\Pi(dV) = \int\limits_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \int\limits_{\{-1,1\}} 1_D(\xi V)\,\lambda(d\xi)\,\omega_d(dV) \quad D \in \mathcal{B}\left(\tilde{\mathbb{S}}_{\mathbb{M}_d}\right),$$

where λ is the spherical measure of ν and ω_d is the probability measure on $\tilde{\mathbb{S}}^+_{\mathbb{M}_d}$ given by (6).

Proof. Let $\lambda(d\xi)$ and ν_ξ be the spherical and radial components of ν given by (4), respectively. For every $z \in \mathbb{R}$, $\mathcal{C}_\mu(z) = i\gamma z + \int_{\mathbb{R}} \left[e^{izx} - 1 - ixz1_{|x|\leq 1}(x)\right]\nu(dx)$ where $\gamma \in \mathbb{R}$ and where we have omitted the Gaussian term without loss of generality. From (5) we have for every Hermitian matrix A,

$$\begin{aligned}
&\log \mathbb{E}[\exp(i\mathrm{tr}(M_d A))] \\
&= d\mathbb{E}_u\left\{i\gamma\langle u, Au\rangle + \int_{\mathbb{R}}\left[e^{i\langle u,Au\rangle x} - 1 - ix\langle u, Au\rangle 1_{|x|\leq 1}(x)\right]\nu(dx)\right\} \\
&= i\gamma r_A d + d\int_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \int_{\mathbb{R}} \left[e^{i\mathrm{tr}(Auu^*)x} - 1 - ix\mathrm{tr}(Auu^*)1_{|x|\leq 1}(x)\right]\nu(dx)\,\omega_d(dV)
\end{aligned}$$

$$= i\gamma r_A d + d \int_{\tilde{\mathbb{S}}^{+}_{\mathbb{M}_d}} \int_{\{-1,1\}} \int_0^\infty \left[e^{i\mathrm{tr}(AV)r\xi} - 1 - ir\xi \mathrm{tr}\left(AV\right) 1_{|r\xi|\leq 1}\left(r\xi\right)\right] \nu_\xi\left(dr\right)\lambda\left(d\xi\right)\omega_d\left(dV\right)$$

$$= i\gamma r_A d + d \int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}} \int_0^\infty e^{i\mathrm{tr}\left(A\tilde{V}\right)r} - 1 - ir\mathrm{tr}\left(A\tilde{V}\right) 1_{r\leq 1}\left(r\right)\nu_\xi\left(dr\right)\Pi\left(d\tilde{V}\right),$$

where $r_A = \mathbb{E}_u \langle u, Au\rangle$ and $\Pi\left(d\tilde{V}\right)$ is a measure on $\tilde{\mathbb{S}}_{\mathbb{M}_d}$ such that for any Borel set D of $\tilde{\mathbb{S}}_{\mathbb{M}_d}$

$$\int\limits_{\left\{\tilde{V}:\ \mathrm{rank}\left(\tilde{V}\right)=1, \left\|\tilde{V}\right\|=1\right\}} 1_D\left(\tilde{V}\right)\Pi\left(d\tilde{V}\right) = \int\limits_{\{V>0:\ \mathrm{rank}(V)=1, \|V\|=1\}} \int\limits_{\{-1,1\}} 1_D\left(\xi V\right)\lambda\left(d\xi\right)\omega_d\left(dV\right),$$

here the equivalence of the regions of integration follows from the spectral representation theorem, since any V with $\mathrm{rank}\left(V\right) = 1$ and $\|V\| = 1$ can be written as $V = \lambda vv^*$ where λ is eigenvalue of V with corresponding eigenvector v, hence $x^*Vx = \lambda\left|x^*v\right|^2$ and therefore $V > 0$ or $V < 0$ according to the sign of λ. ■

3 Random matrix models of stochastic integral type

Any real-valued Lévy process $\left\{X_t^{(\mu)} : t \geq 0\right\}$ with law μ at time $t = 1$, uniquely induces a real-valued independently scattered random measure $\left\{M^{(\mu)}\left(B\right) : B \in \mathcal{B}_0\left([0,\infty)\right)\right\}$ such that $M^{(\mu)}\left([0,t]\right) = X_t^{(\mu)}$ almost surely, where $\mathcal{B}_0\left([0,\infty)\right)$ is the family of bounded Borel sets in $[0,\infty)$. We will consider $M^{(\mu)}$-integrable $\left(\text{or } X_t^{(\mu)}\text{-integrable}\right)$ real-valued functions h on $[0,\infty)$ in the sense of [15] and [19]. Then $\int_B h(t)M^{(\mu)}(dt)$ $\left(\text{or } \int_B h(t)dX_t^{(\mu)}\right)$ is defined almost surely for every $B \in \mathcal{B}_0\left([0,\infty)\right)$. The stochastic integral of a real-valued h on $[0,\infty)$ with respect to $X_t^{(\mu)}$ is a real-valued infinitely divisible random variable written as

$$\eta = \int_0^\infty h(t)dX_t^{(\mu)}, \tag{7}$$

which is defined as the limit in probability of $\int_{[0,s]} h(t)dX_t^{(\mu)}$ as $s \to \infty$ whenever the limit exists. Furthermore, its cumulant transform is given by

$$\mathcal{C}_\eta\left(z\right) = \int_0^\infty \mathcal{C}_\mu\left(h(t)z\right)dt \quad z \in \mathbb{R}. \tag{8}$$

For the complex matrix case we have a similar result; see [5] for the case of real matrices. For any infinitely divisible matrix Ψ in $\mathbb{M}_d$ with associated matrix Lévy process $\left\{\Psi_t^d : t \geq 0\right\}$, the infinitely divisible $d \times d$ matrix valued stochastic integral

$$M = \int_0^\infty h(t)d\Psi_t^d, \tag{9}$$

whenever exists, has cumulant transform

$$\mathcal{C}_M(A) = \int_0^\infty \mathcal{C}_\Psi(h(t)A)\,dt \quad A \in \mathbb{M}_d. \tag{10}$$

In this work we consider matrix Lévy processes $\left\{\Psi_t^d\right\}$ corresponding to Lévy measures of the form

$$\nu_\Psi^d(B) = d\int_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \omega_d(dV) \int_{\mathbb{R}} 1_B(xV)\,\nu_\mu(dx), \tag{11}$$

where ω_d is the probability measure induced by the transformation $u \to V = uu^*$ in (6) and ν_μ is a Lévy measure of an infinitely divisible distribution μ on $\mathbb{R}$. Observe that ν_Ψ^d is Lévy measure supported in the subset of rank one matrices in $\mathbb{M}_d$.

The following is the main result of this work, it provides random matrix models for free infinitely divisible distributions on $\mathbb{R}$ given by matrix stochastic integrals of the form (9) when the corresponding classical infinitely divisible distributions under Λ are representable as the random integrals of the form (7).

Theorem 2 *Let μ_h be an infinitely divisible distribution on $\mathbb{R}$ given by the stochastic integral representation*

$$\mu_h = \mathcal{L}\left(\int_0^\infty h(t)dX_t^{(\mu)}\right), \tag{12}$$

where $X_t^{(\mu)}$ is a Lévy process on $\mathbb{R}$ with law μ at time $t = 1$ and Lévy measure ν_μ. The free infinitely divisible distribution $\Lambda(\mu_h)$ has a random matrix model given by the ensemble of infinitely divisible matrix stochastic integrals

$$\left(M_h^d = \int_0^\infty h(t)d\Psi_t^d\right)_{d\geq 1}, \tag{13}$$

where Ψ_t^d is the $\mathbb{M}_d$-valued Lévy process with Lévy measure ν_Ψ^d given by (11) in terms of ω_d and ν_μ.

Proof. We will proof that the random matrices of the ensemble $\left(M_h^d\right)_{d\geq 1}$ in (13) and the random matrices of the random matrix model $(M_d)_{d\geq 1}$ for $\Lambda(\mu_h)$ given in [6, Theorem 6.1] have the same laws. Recall from (5) that the Fourier transform of M_d is given by

$$\mathbb{E}[\exp(i\mathrm{tr}(M_d A))] = \exp\left[d\mathbb{E}_u \mathcal{C}_{\mu_h}(\langle u, Au\rangle)\right] \ \ A \text{ Hermitian}.$$

We will prove that this Fourier transform of M_d coincides with the Fourier transform of M_h^d.

For that, we first calculate the cumulant transform of Ψ_t^d at time 1 using (11),

$$\begin{aligned}
\mathcal{C}_{\Psi_1^d}(A) &= \int_{\mathbb{M}_d} \left[e^{i\mathrm{tr}(AX)} - 1 - i\mathrm{tr}(AX) 1_{\|X\|\leq 1}(X) \right] \nu_{\Psi_1}^d(dX) \\
&= d \int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}^+} \int_{\mathbb{R}} \left[e^{i\mathrm{tr}(AV)x} - 1 - ix\mathrm{tr}(AV) 1_{|x|\leq 1}(x) \right] \omega_d(dV)\, \nu_\mu(dx) \\
&= d \int_{\mathbb{R}} \mathbb{E}_V \left[e^{i\mathrm{tr}(AV)x} - 1 - ix\mathrm{tr}(AV) 1_{|x|\leq 1}(x) \right] \nu_\mu(dx) \\
&= d\mathbb{E}_u \int_{\mathbb{R}} \left[e^{i\mathrm{tr}(Auu^*)x} - 1 - ix\mathrm{tr}(Auu^*) 1_{|x|\leq 1}(x) \right] \nu_\mu(dx) \\
&= d\mathbb{E}_u \mathcal{C}_\mu(\langle u, Au \rangle). \qquad (14)
\end{aligned}$$

Now we calculate the cumulant transform of M_h^d from (10) and (14),

$$\begin{aligned}
\mathcal{C}_{M_h^d}(A) &= \int_0^\infty \mathcal{C}_{\Psi_1^d}(h(t)A)\, dt = \int_0^\infty d\mathbb{E}_u \mathcal{C}_\mu(\langle u, h(t)Au \rangle)\, dt \\
&= d\mathbb{E}_u \int_0^\infty \mathcal{C}_\mu(h(t)\langle u, Au \rangle)\, dt = d\mathbb{E}_u \mathcal{C}_{\mu_h}(\langle u, Au \rangle),
\end{aligned}$$

where in the last equality we have used the relation (8) between the cumulant transforms of μ_h and μ corresponding to the stochastic integral representation (12). ■

Remark 3 *Let $\nu_\mu, \nu_{\mu_h}, \nu_{M_h^d}$ and ν_{Ψ^d} denote the Lévy measures of μ, μ_h, M_h^d and Ψ_t^d at time 1 in Theorem 2, respectively. From [19] the Lévy measures of μ and μ_h are related as follows*

$$\nu_{\mu_h}(B) = \int_0^\infty dt \int_{\mathbb{R}} 1_B(h(t)x)\nu_\mu(dx) \quad B \in \mathcal{B}(\mathbb{R}\backslash\{0\})$$

and from [5] it is obtained a similar relation between the Lévy measures of M_h^d and Ψ^d,

$$\nu_{M_h^d}(B) = \int_0^\infty dt \int_{\mathbb{M}_d} 1_B(h(t)X)\nu_{\Psi^d}(dX) \quad B \in \mathcal{B}(\mathbb{M}_d\backslash\{0\}).$$

These two relations combined with (11) yield the Lévy measure of M_h^d in terms of the Lévy measure of μ_h

$$\begin{aligned}
\nu_{M_h^d}(B) &= d \int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}^+} \omega_d(dV) \int_{\mathbb{R}} \nu_\mu(dx) \int_0^\infty 1_B(h(t)xV) dt \\
&= d \int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}^+} \omega_d(dV) \int_{\mathbb{R}} \nu_{\mu_h}(dx)\, 1_B(xV), \quad B \in \mathcal{B}(\mathbb{M}_d\backslash\{0\}). \qquad (15)
\end{aligned}$$

4 Examples

In the following $I(\mathbb{R})$ denotes the class of infinitely divisible distributions on $\mathbb{R}$ and $L(\mathbb{R})$ denotes the class of selfdecomposable distributions on $\mathbb{R}$.

Let (λ, ν_ξ) denote the polar decomposition of the Lévy measure ν of any $\mu \in I(\mathbb{R})$ given by (4), that is

$$\nu(B) = \int_S \lambda(d\xi) \int_0^\infty 1_B(r\xi)\nu_\xi(dr),$$

where λ and ν_ξ are the spherical and radial components of ν, respectively and $S = \{-1, 1\}$. Moreover, let (λ, k_ξ) denote the following description of the above descomposition for selfdecomposable distributions, see [18]. If $\mu \in L(\mathbb{R})$ the radial component of its Lévy measure ν is expressed as

$$\nu_\xi(dr) = 1_{(0,\infty)}(r)\frac{k_\xi(r)}{r}dr,$$

where $k_\xi(r)$ is a nonnegative measurable function in $\xi \in S$ and decreasing, right continuous in $r \in (0, \infty)$. Here k_ξ is called the k-function of ν.

Throughout this section we provide examples of random matrix models for free infinitely divisible distributions $\Lambda(\mu_h)$ given by the ensembles of matrix stochastic integrals (13), for several classes infinitely divisible distributions μ_h with stochastic integral representation (12). In most of examples we express the Lévy measure $\nu_{M_h^d}$ in terms of ν_{μ_h}. We emphasize, according to (15), that these Lévy measures $\nu_{M_h^d}$ are supported in the subset of rank one matrices in $\mathbb{M}_d$.

In all examples below $\nu_\mu, \nu_{\mu_h}, \nu_{M_h^d}$ and ν_{Ψ^d} denote the Lévy measures of μ, μ_h, M_h^d and Ψ_t^d at time 1 in Theorem 2, respectively.

The following two examples recover the random matrix models of Theorems 4.1 and 4.3 in [14] and find their corresponding Lévy measures.

Example 1. The class of Generalized Gamma Convolutions $T(\mathbb{R}_+)$ is the smallest class of infinitely divisible distributions on $\mathbb{R}_+$ that contains all Gamma distributions and that is closed under convolution and weak convergence.

In [9], any distribution in $T(\mathbb{R}_+)$ has stochastic integral representation (12) where μ is a Gamma distribution and $h(t)$ is a Borel measurable function $h : \mathbb{R}_+ \to \mathbb{R}_+$ such that $\int_0^\infty \log(1 + h(t))\, dt < \infty$. Such a representation is called the Wiener-Gamma integral representation.

Let $\mu_h \in T(\mathbb{R}_+)$ has the Wiener-Gamma integral representation. By (8) in [14] the Lévy measure of μ_h is expressed as

$$\nu_{\mu_h}(dx) = 1_{(0,\infty)}(x) \int_0^\infty \frac{e^{-x/h(t)}}{x} dt dx.$$

The corresponding random matrix models (13) for the free Generalized Gamma Convolutions $\Lambda(\mu_h)$ consist of matrix stochastic integrals of Wiener-Gamma type

$$\left(M_h^d = \int_0^\infty h(t) d\Psi_t^d \right)_{d \geq 1}$$

with Lévy measures

$$\nu_{M_h^d}(B) = d \int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}^+} \omega_d\,(dV) \int_0^\infty dx 1_B(xV) \int_0^\infty \frac{e^{-x/h(t)}}{x} dt.$$

Example 2. The class of Thorin distributions $T(\mathbb{R})$ is the smallest class of infinitely divisible distributions on $\mathbb{R}$ which contains all distributions in $T(\mathbb{R}_+)$ and is closed under convolution, weak convergence and reflection.

It is shown in [4] that this class is characterized by the stochastic integral representation (12) where $\mu \in ID_{\log}(\mathbb{R})$ and $h(t)$ is the inverse function of the incomplete Gamma function $g(t) = \int_t^\infty e^{-s} s^{-1} ds$.

Let $\mu_h \in T(\mathbb{R})$ with such a representation. The random matrix models (13) for the corresponding free Thorin distributions $\Lambda(\mu_h)$ are given by

$$\left(M_h^d = \int_0^\infty g^*(t) d\Psi_t^d \right)_{d\geq 1} \qquad g^* \text{ inverse function of } g.$$

Let (λ, ν_ξ) and (λ_h, ν_{h_ξ}) be the polar decompositions of the Lévy measures of μ and μ_h, respectively. It is proved in [1] that the corresponding k-function of μ_h is $k_{h_\xi}(r) = \int_0^\infty e^{-r/s} \nu_\xi(ds)$. Therefore the Lévy measures of these matrix stochastic integrals are given by

$$\nu_{M_h^d}(B) = d \int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}^+} \omega_d\,(dV) \int_S \lambda_h(d\xi) \int_0^\infty \frac{dr}{r} \left(\int_0^\infty e^{-r/s} \nu_\xi(ds) \right) 1_B(r\xi V).$$

Example 3. The class of Bondesson distributions $B(\mathbb{R})$ is the smallest class of infinitely divisible distributions on $\mathbb{R}$ that contains all mixtures of exponential distributions and that is closed under convolution, weak convergence and reflection.

This class is characterized by the stochastic integral representation (12) where $\mu \in I(\mathbb{R})$ and $h(t) = 1_{(0,1)}(t) \log(1/t)$, see [4]. If $\mu_h \in B(\mathbb{R})$ has such a representation then by (2.17) in [4]

$$\nu_{\mu_h}(dx) = \int_0^\infty e^{-s} \nu_\mu \left(s^{-1} dx \right) ds.$$

Therefore the random matrix models (13) for the free Bondesson distributions $\Lambda(\mu_h)$ are given by

$$\left(M_h^d = \int_0^1 \log(1/t)\, d\Psi_t^d \right)_{d\geq 1}$$

with Lévy measures

$$\nu_{M_h^d}(B) = d \int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}^+} \omega_d\,(dV) \int_{-\infty}^\infty \int_0^\infty e^{-s} \nu_\mu \left(s^{-1} dx \right) ds 1_B(xV).$$

Example 4. The class of Thorin distributions $T(\mathbb{R})$ is also characterized (see Example 2) by the stochastic integral representation (12) where $\mu \in L(\mathbb{R})$ and $h(t) = 1_{(0,1)}(t) \log(1/t)$,

see [4].
If $\mu_h \in T(\mathbb{R})$ has this alternative representation, the corresponding random matrix models (13) for the free Thorin distributions $\Lambda(\mu_h)$ are

$$\left(M_h^d = \int_0^1 \log(1/t)\, d\Psi_t^d \right)_{d\geq 1}.$$

Let (λ, k_ξ) and (λ_h, k_{h_ξ}) be the corresponding polar decompositions of the Lévy measures of μ and μ_h, respectively. It is shown in [4] that $\lambda = \lambda_h$ and

$$k_{h_\xi}(r) = \int_0^\infty k_\xi(rs^{-1}) e^{-s} ds$$

and hence the Lévy measures for these random matrix models are of the form

$$\begin{aligned} \nu_{M_h^d}(B) &= d \int_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \omega_d\,(dV) \int_{-\infty}^\infty 1_B(xV) \nu_{\mu_h}(dx) \\ &= d \int_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \omega_d\,(dV) \int_S \lambda\,(d\xi) \int_0^\infty \frac{dr}{r} \left(\int_0^\infty k_\xi(rs^{-1}) e^{-s} ds \right) 1_B(r\xi V). \end{aligned}$$

Example 5. The class of selfdecomposable distributions $L(\mathbb{R})$ is characterized by the stochastic integral representation (12) where $\mu \in I_{\log}(\mathbb{R})$ and $h(t) = 1_{(0,\infty)}(t)e^{-t}$, see [11], [20] and [18].

Let $\mu_h \in L(\mathbb{R})$ with such a Ornstein-Uhlenbeck type integral representation. The random matrix models (13) for the corresponding free selfdecomposable distributions $\Lambda(\mu_h)$ are given by

$$\left(M_h^d = \int_0^\infty e^{-t} d\Psi_t^d \right)_{d\geq 1}$$

and satisfy an $I_{\log}$-condition, that is

$$\int_{\|X\|>2} \log\|X\|\, \nu_\Psi^d(dX) < \infty,$$

which follows easily from (11) and the fact that $\mu \in I_{\log}(\mathbb{R})$. These Ornstein-Uhlenbeck type matrix integrals are supported on the subset of rank one matrices in $\mathbb{M}_d$.
Let (λ, ν_ξ) and (λ_h, k_{h_ξ}) be the corresponding polar decompositions of the Lévy measures of μ and μ_h. It is proved in [1] that the k-function of ν_{μ_h} is given by

$$k_{h_\xi}(r) = \nu_\xi\left((r,\infty)\right).$$

Therefore, the Lévy measures of these random matrix models of Ornstein-Uhlenbeck type for free selfdecomposable distributions $\Lambda(\mu_h)$ are given by

$$\begin{aligned} \nu_{M_h^d}(B) &= d \int_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \omega_d\,(dV) \int_{-\infty}^\infty 1_B(xV) \nu_{\mu_h}(dx) \\ &= d \int_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \omega_d\,(dV) \int_S \lambda_h\,(d\xi) \int_0^\infty \frac{dr}{r} \nu_\xi\left((r,\infty)\right) 1_B(r\xi V). \end{aligned}$$

Example 6. The class of type G distributions $G_{sym}(\mathbb{R})$ is the class of symmetric distributions on $\mathbb{R}$ which are variance mixtures of the standard Gaussian distribution with positive infinitely divisible mixing distributions.

In [2] is proved that this class is characterized by the stochastic integral representation (12) where $\mu \in I(\mathbb{R})$ and $h(t)$ is the inverse function of $f(t) = \int_t^\infty \varphi(u)du$ where $\varphi(u) = \frac{1}{\sqrt{2\pi}} e^{-\frac{u^2}{2}}$.

Let $\mu_h \in G_{sym}(\mathbb{R})$ with such a representation. In this case the random matrix models (13) for the corresponding free type G distributions $\Lambda(\mu_h)$ are of the form

$$\left(M_h^d = \int_0^\infty f^*(t) d\Psi_t^d \right)_{d \geq 1} \qquad f^* \text{ inverse function of } f.$$

Let (λ_h, ν_{h_ξ}) be the polar decomposition of the Lévy measure of the type G distribution μ_h. It is known that the radial component ν_{h_ξ} can be written as

$$\nu_{h_\xi}(dr) = g_{h_\xi}(r^2) dr$$

where $g_{h_\xi}(r)$ is measurable in $\xi \in S$ and completely monotone in $r \in (0, \infty)$. It is proved in [1] that

$$g_{h_\xi}(r) = \int_0^\infty \varphi(r^{1/2}/s) s^{-1} \nu_\xi(ds),$$

where ν_ξ is the radial component of ν_μ. Therefore the Lévy measures for these random matrix models for free type G distributions $\Lambda(\mu_h)$ are given by

$$\nu_{M_h^d}(B) = d \int_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \omega_d(dV) \int_S \lambda_h(d\xi) \int_0^\infty dr \left(\int_0^\infty \varphi(r/s) s^{-1} \nu_\xi(ds) \right) 1_B(r\xi V).$$

Example 7. The class of M distributions $M(\mathbb{R})$ is a subclass of selfdecomposable distributions of $G_{sym}(\mathbb{R})$. It is the class of symmetric infinitely divisible distributions on $\mathbb{R}$ whose Lévy measures have polar decomposition (λ, ν_ξ) such that

$$\nu_\xi(dr) = \frac{g_\xi(r^2)}{r} dr, \tag{16}$$

where $g_{h_\xi}(r)$ is measurable in $\xi \in S$ and completely monotone in $r \in (0, \infty)$.

In [3] is shown that the class $M(\mathbb{R})$ is characterized by the stochastic integral representation (12) where $\mu \in I_{\log}(\mathbb{R})$ and $h(t)$ is the inverse function of $m(t) = \int_t^\infty \frac{\varphi(u)}{u} du$.

Let $\mu_h \in M(\mathbb{R})$ with such a representation. The random matrix models (13) for the corresponding free M distributions $\Lambda(\mu_h)$ are

$$\left(M_h^d = \int_0^\infty m^*(t) d\Psi_t^d \right)_{d \geq 1} \qquad m^* \text{ inverse function of } m.$$

Let (λ_h, ν_{h_ξ}) be the polar decomposition of the Lévy measure of μ_h and let g_{h_ξ} be the corresponding g-function in (16) of the radial component ν_{h_ξ}. It is proved in [1] that

$$g_{h_\xi}(r) = \int_0^\infty \varphi(r^{1/2}/s)\nu_\xi(ds),$$

where ν_ξ is the radial component of ν_μ. Therefore the Lévy measures for these random matrix models for the free M distributions $\Lambda(\mu_h)$ can written as

$$\nu_{M_h^d}(B) = d\int_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \omega_d\,(dV) \int_S \lambda_h\,(d\xi) \int_0^\infty dr \left(\int_0^\infty \varphi(r/s)\nu_\xi(ds) \right) 1_B(r\xi V).$$

Example 8. The class of Jurek distributions $U(\mathbb{R})$ is the class of infinitely divisible distributions on $\mathbb{R}$ whose Lévy measures have polar decomposition (λ, ν_ξ) such that

$$\nu_\xi\,(dr) = l_\xi(r)dr, \tag{17}$$

where l_ξ is measurable in $\xi \in S$ and decreasing in $r \in (0, \infty)$.

It is shown in [10] that this class is characterized by the stochastic integral representation (12) where $\mu \in I(\mathbb{R})$ and $h(t) = 1_{[0,1]}(t)t$.

Let $\mu_h \in U(\mathbb{R})$ with such a representation. The corresponding random matrix models (13) for the free Jurek distributions $\Lambda(\mu_h)$ are of the form

$$\left(M_h^d = \int_0^1 t d\Psi_t^d \right)_{d\geq 1}.$$

Let (λ, ν_ξ) and (λ_h, ν_{h_ξ}) be the polar decompositions of the Lévy measures of μ and μ_h, respectively. It is proved in [1] that the corresponding l-function in (17) for the radial component ν_{h_ξ} is expressed as $l_{h_\xi}(r) = \int_r^\infty x^{-1}\nu_\xi(dx)$. Therefore the Lévy measures of these random matrix models are expressed as

$$\nu_{M_h^d}(B) = d\int_{\tilde{\mathbb{S}}^+_{\mathbb{M}_d}} \omega_d\,(dV) \int_S \lambda_h\,(d\xi) \int_0^\infty dr \left(\int_r^\infty x^{-1}\nu_\xi(dx) \right) 1_B(r\xi V).$$

Example 9. The class of A distributions $A(\mathbb{R})$ introduced in [12] is bigger than the Jurek class $U(\mathbb{R})$. It is the class of infinitely divisible distributions on $\mathbb{R}$ with Lévy measures ν of the form

$$\nu(B) = \int_{\mathbb{R}\backslash\{0\}} \rho\,(dx) \int_0^\infty a_1\,(r; |x|)\, 1_B \left(r\frac{x}{|x|} \right) dr \qquad B \in \mathcal{B}\,(\mathbb{R})\,,$$

where ρ is a Lévy measure on $\mathbb{R}$; and $a_1\,(r; s)$ is the one-sided arcsine density with parameter $s > 0$, that is,

$$a_1\,(r; s) = \begin{cases} 2\pi^{-1}\,(s - r^2)^{-1/2} & 0 < r < s^{1/2} \\ 0 & \text{otherwise.} \end{cases}$$

It is shown in [12] that the class $A(\mathbb{R})$ is characterized by the stochastic integral representation (12) where $\mu \in I(\mathbb{R})$ and $h(t) = 1_{[0,1]}(t)\cos\left(\frac{\pi}{2}t\right)$.

Let $\mu_h \in A(\mathbb{R})$ with such a representation. In this case the random matrix models (13) for the corresponding free A distributions $\Lambda(\mu_h)$ are

$$\left(M_h^d = \int_0^1 \cos\left(\frac{\pi}{2}t\right) d\Psi_t^d\right)_{d\geq 1}$$

with Lévy measures, see (15),

$$\nu_{M_h^d}(B) = d\int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}^+} \omega_d(dV) \int_{\mathbb{R}} \nu_\mu(dx) \int_0^1 1_B\left(xV\cos\left(\frac{\pi}{2}t\right)\right) dt.$$

Example 10. The class of Type G distributions $G_{sym}(\mathbb{R})$ is also characterized (see Example 6) by the stochastic integral representation (12) where $\mu \in A(\mathbb{R})$ and $h(t) = 1_{(0,1/2)}(t)\left(\log\frac{1}{t}\right)^{1/2}$, see [12].
Let $\mu_h \in G_{sym}(\mathbb{R})$ has this alternative representation, that is

$$\mu_h = \mathcal{L}\left(\int_0^{1/2}\left(\log\frac{1}{t}\right)^{1/2} dX_t^{(\mu)}\right) \quad \mu \in A(\mathbb{R}). \tag{18}$$

The corresponding random matrix models (13) for the free Type G distributions $\Lambda(\mu_h)$ are given by

$$\left(M_h^d = \int_0^{1/2}\left(\log\frac{1}{t}\right)^{1/2} d\Psi_t^d\right)_{d\geq 1}$$

with Lévy measures

$$\nu_{M_h^d}(B) = d\int_{\tilde{\mathbb{S}}_{\mathbb{M}_d}^+} \omega_d(dV) \int_{\mathbb{R}} \nu_\mu(dx) \int_0^{1/2} 1_B\left(xV\left(\log\frac{1}{t}\right)^{1/2}\right) dt.$$

Example 11. The class of generalized Type G distributions $G(\mathbb{R})$ introduced in [13] is the class of infinitely divisible distributions on $\mathbb{R}$ whose Lévy measures have polar decomposition (λ, ν_ξ) such that

$$\nu_\xi(dr) = g_\xi(r^2)dr,$$

where $g_\xi(r)$ is measurable in $\xi \in S$ and completely monotone in $r \in (0,\infty)$. Distributions of this class are not necessarily symmetric. If $\mu \in G(\mathbb{R})$ is symmetric then $\mu \in G_{sym}(\mathbb{R})$, that is μ is of Type G.

It is proved in [12] that the class of generalized type G distributions is characterized by the stochastic integral representation (12) where $\mu \in A(\mathbb{R})$ and $h(t) = 1_{(0,1)}(t)\left(\log\frac{1}{t}\right)^{1/2}$.

Let $\mu_h \in G(\mathbb{R})$ has such a representation, that is

$$\mu_h = \mathcal{L}\left(\int_0^1\left(\log\frac{1}{t}\right)^{1/2} dX_t^{(\mu)}\right) \quad \mu \in A(\mathbb{R}). \tag{19}$$

This representation (19) is not necessarily a symmetric generalization of (18).

In this case the corresponding random matrix models (13) for the free generalized type G distributions $\Lambda(\mu_h)$ are

$$\left(M_h^d = \int_0^1 \left(\log \frac{1}{t} \right)^{1/2} d\Psi_t^d \right)_{d \geq 1}.$$

References

[1] T. Aoyama and M. Maejima: Some classes of infinitely divisible distributions on $\mathbb{R}^d$ (a survey), *unpublished note* (2008).

Revised version of the Research Report: T. Aoyama and M. Maejima: Some classes of infinitely divisible distributions on $\mathbb{R}^d$ (a survey), *The Institute of Statistical Mathematics Cooperate Research Report*, Tokyo, **184**, (2006), 5-13.

[2] T. Aoyama and M. Maejima: Characterizations of subclasses of type G distributions on $\mathbb{R}^d$ by stochastic integral representations, *Bermoulli*, **13**(1), (2007), 148-160.

[3] T. Aoyama, M. Maejima and J. Rosiński: A subclass of type G selfdecomposable distributions, *J. Theoret. Probab.*, **21**, (2008), 14-34.

[4] O.E. Barndorff-Nielsen, M. Maejima and K. Sato: Some classes of multivariate infinitely divisible distributions admitting stochastic integral representations, *Bernoulli,* **12**, (2006), 1–33.

[5] O. E. Barndorff-Nielsen and R. Stelzer: Multivariate supOU processes, Ann. Appl. Probab., 21 (2011), 140-182.

[6] F. Benaych-Georges: Classical and free infinitely divisible distributions and random matrices, *Ann. Probab.,* **33**, (2005), 1134–1170.

[7] H. Bercovici and V. Pata: Stable laws and domains of attraction in free probability theory, *Ann. of Math.*, **149,** (1999), 1023-1060.

[8] T. Cabanal-Duvillard: A matrix representation of the Bercovici-Pata bijection, *Electron. J. Probab.*, **10** (2005), 632-661.

[9] L. F. James, B. Roynette and M. Yor: Generalized Gamma convolutions, Dirichlet means, Thorin measures, with explicit examples, *Probability Surveys*, **5**, (2008), 346-415.

[10] Z. J. Jurek: Relations between the s-selfdecomposable and selfdecomposable measures, *Ann. Probab.*, **13**, (1985), 592-608.

[11] Z. J. Jurek and W. Vervaat: An integral representation for selfdecomposable Banach space valued random variables, *Z. Wahrscheinlichkeitstheorie. Verw. Geb.*, **62,** (1983), 247-262.

[12] M. Maejima, V. Pérez-Abreu and K. Sato: A class of multivariate infinitely divisible distributions related to arcsine density, Bernoulli (to appear).

[13] M. Maejima and K. Sato: The limits of nested subclasses of several classes of infinitely divisible distributions are identical with the closure of the class of stable distributions, *Probab. Theory Relat. Fields*, **145**, (2009), 119-142.

[14] V. Pérez-Abreu and N. Sakuma: Free generalized gamma convolutions. *Elect. Comm. in Probab.*, **13**, (2008), 526-539.

[15] B. S. Rajput and J. Rosiński: Spectral representations of infinitely divisible processes, *Probab. Theory Related Fields*, **82,** (1989), 451–487.

[16] A. Rocha-Arteaga and K. Sato: *Topics in Infinitely Divisible Distributions and Lévy Processes*, Aportaciones Mat., Invest., **17,** (2003) México: Sociedad Matemática Mexicana.

[17] J. Rosiński: On series representations of infinitely divisible random vectors, *Ann. Probab.*, **18,** (1990), 405-430.

[18] K. Sato: *Lévy Processes and Infinitely Divisible Distributions*, Cambridge Univ. Press, Cambridge, (1999).

[19] K. Sato: Additive Processes and Stochastic Integrals. *Illinois J. of Math.*, **50,** (2006), 825-851.

[20] K. Sato and M. Yamazato: Stationary processes of Ornstein-Uhlenbeck type, In K. Ito and J. V. Prokhorov (eds), *Probability Theory and Mathematical Statistics*, Lectures. Notes in Math. **1021**, Berlin: Spriger-Verlag, (1983), 541-551.

[21] K. Urbanik and W.A. Woyczynski: A random integral and Orlicz spaces. *Bull. Acad. Polon. Sci. Math. Astro. e Phys.*, **15**, (1967), 161–168.